



Beckfoot

Subject: Biology

Topic: Biological molecules (I)

Year Group: I2


enjoy
learn
succeed

Key Vocabulary

| | | |
|----|-----------------------|---|
| 1 | Monomer | Individual molecules that make up a polymer |
| 2 | Polymer | Long chains composed of many individual monomers bonded together in repeating pattern |
| 3 | Condensation reaction | Occurs when two molecules combine to form a more complex molecule with the removal of water |
| 4 | Hydrolysis reaction | Occurs when larger molecules are broken down into smaller molecules with the addition of water |
| 5 | Monosaccharide | Simplest carbohydrates, consisting of only one sugar molecule eg: glucose, fructose, galactose |
| 6 | Disaccharide | Sugars composed of two monosaccharides joined together by a glycosidic bond, in a condensation reaction eg: maltose, sucrose, lactose. Can be separated by a hydrolysis reaction. |
| 7 | Polysaccharide | Sugars composed of many monosaccharides joined together by glycosidic bonds, in a condensation reaction eg: amylose, glycogen, cellulose. Can be separated by a hydrolysis reaction. |
| 8 | Amino acid | Monomer units that make up proteins. 20 amino acids exist with different R groups |
| 9 | Dipeptide | Two amino acids joined together by a peptide bond in a condensation reaction. Can be separated by a hydrolysis reaction. |
| 10 | Polypeptide | A polymer made of many amino acids joined together by peptide bonds in a condensation reaction. Can be separated by a hydrolysis reaction. |
| 11 | Protein | Contains one or more polypeptide chains. There are four structural levels |

Monomers and polymers

| Monomers and polymers (see key vocabulary table) | | |
|--|---------------------------------------|--|
| 1 | Condensation reaction (see key vocab) | |
| 2 | Hydrolysis reaction (see key vocab) | |







Carbohydrates

| | | | |
|---|---|--|--|
| 1 | Monosaccharides Eg: glucose, fructose, galactose (see key vocab) | Glucose is a hexose sugar with 2 isomers | |
| 2 | Disaccharides (see key vocab) Eg: maltose, sucrose, lactose | Name | Monosaccharide constituent |
| | | a) Maltose | 2 x α -glucose |
| | | b) Sucrose | α -glucose and fructose |
| | | c) Lactose | β -glucose and galactose |
| 3 | Polysaccharides (see key vocab) Eg: Amylose, glycogen, cellulose | Name | Function |
| | | a) Amylose | Amylopectin (starch) is the main polysaccharide energy store in plants. It is composed of alpha glucose. |
| | | b) Glycogen | Polysaccharide energy store in animals, composed of alpha glucose |
| | | c) Cellulose | Structural component of plant cell walls, composed of long unbranched chains of beta glucose |

Proteins

| 1 | Amino acids (see key vocab) | | 4 | Protein (see key vocab) | | | | | | |
|------------|---|---|---|--|------------|---|---|---------|---|---------------|
| 2 | Dipeptide (see key vocab) | <p>A dipeptide</p> | 5 | <table><tr><th>Level</th><th>Definition</th><th>Bond type</th></tr><tr><td>Primary</td><td>The specific sequence of amino acids in a polypeptide chain</td><td>Peptide bonds</td></tr></table> | Level | Definition | Bond type | Primary | The specific sequence of amino acids in a polypeptide chain | Peptide bonds |
| Level | Definition | Bond type | | | | | | | | |
| Primary | The specific sequence of amino acids in a polypeptide chain | Peptide bonds | | | | | | | | |
| 3 | Polypeptide (see key vocab) | | 6 | <table><tr><td>Secondary</td><td>The curling or folding of the polypeptide chain into α-helices and β-pleated sheets due to the formation of hydrogen bonds</td><td>Hydrogen bonds</td></tr></table> | Secondary | The curling or folding of the polypeptide chain into α -helices and β -pleated sheets due to the formation of hydrogen bonds | Hydrogen bonds | | | |
| Secondary | The curling or folding of the polypeptide chain into α -helices and β -pleated sheets due to the formation of hydrogen bonds | Hydrogen bonds | | | | | | | | |
| | | | 7 | <table><tr><td>Tertiary</td><td>The overall specific 3-D shape of a protein, which is determined by interactions between R groups and the properties of R groups</td><td>Hydrogen bonds Ionic bonds Disulphide bridges</td></tr></table> | Tertiary | The overall specific 3-D shape of a protein, which is determined by interactions between R groups and the properties of R groups | Hydrogen bonds Ionic bonds Disulphide bridges | | | |
| Tertiary | The overall specific 3-D shape of a protein, which is determined by interactions between R groups and the properties of R groups | Hydrogen bonds Ionic bonds Disulphide bridges | | | | | | | | |
| | | | 8 | <table><tr><td>Quaternary</td><td>The specific 3-D shape of a protein that is determined by the multiple polypeptide chains and/or prosthetic groups bonded together</td><td>Hydrogen bonds Ionic bonds Disulphide bridges</td></tr></table> | Quaternary | The specific 3-D shape of a protein that is determined by the multiple polypeptide chains and/or prosthetic groups bonded together | Hydrogen bonds Ionic bonds Disulphide bridges | | | |
| Quaternary | The specific 3-D shape of a protein that is determined by the multiple polypeptide chains and/or prosthetic groups bonded together | Hydrogen bonds Ionic bonds Disulphide bridges | | | | | | | | |

Lipids



| | | | |
|---|---------------------------------|--|---|
| 1 | Triglyceride (see key vocab) | Structure | Function |
| | |  | Roles in respiration, energy storage- due to insolubility and high carbon to hydrogen ratio |
| 2 | Phospholipid (see key vocab) |  | Hydrophobic tails (water hating) and hydrophilic heads (water loving) allow the phospholipids to form phospholipid bilayers (membranes) |
| 3 | Saturated (see key vocab) | <p>Saturated Lipid</p>  <p>● = Carbon atom ○ = Hydrogen atom</p> | |
| 4 | Unsaturated (see key vocab) | <p>Unsaturated Lipid</p>  <p>● = Carbon atom ○ = Hydrogen atom</p> | |

Biochemical (food) tests

| | | | |
|---|--|---|------------------------|
| 1 | Biochemical tests are use reagents to distinguish the biological molecules contained in food samples | | |
| | Molecule | Reagent | Positive result |
| 2 | Reducing sugars | Benedict's reagent → Heat | Red/orange precipitate |
| 3 | Starch | Iodine in potassium iodide solution | Blue/black |
| 4 | Non-reducing sugars | Hydrochloric acid → Heat Sodium hydrogencarbonate Benedict's reagent → Heat | Red/orange precipitate |
| 5 | Proteins | Sodium hydroxide Copper (II) sulphate | Purple |
| 6 | Lipids | Ethanol Water → Shake | Cloudy white |

Enzymes (1)

There are two models proposed for how enzymes (see key vocab) function. The specificity of enzymes is due to the tertiary structure of its active site, allowing complementary binding to substances

| | |
|---|--|
| 1 | Lock and key model |
| |  <p>enzyme + substrate entering active site → enzyme/substrate complex → enzyme/product complex → enzyme + product leaving active site</p> |
| 2 | Induced fit model |
| |  <p>Substrate entering active site of enzyme → Enzyme/substrate complex → Enzyme/products complex → Products leaving active site of enzyme</p> |


Key Vocabulary

| | | |
|---|--------------------|--|
| 1 | Lipids | Macromolecules made of fatty acid monomers. 2 types: triglycerides, phospholipids |
| 2 | Saturated | Molecule contains no carbon double bonds (C=C), only has carbon single bonds (C-C) has as many hydrogen atoms as possible |
| 3 | Unsaturated | Molecule contains at least one C=C bond and has fewer hydrogen atoms than is maximally possible |
| 4 | Triglyceride | Molecule formed by the joining of one glycerol to three fatty acids by a ester bonds, through condensation reactions. 3 molecules of water are produced as 3 reactions occur. Can be separated by a hydrolysis reaction. |
| 5 | Phospholipid | Molecule formed by the joining of one glycerol to two fatty acids and one phosphate molecule by ester bonds, through condensation reactions |
| 6 | Enzymes | Biological catalysts that speed up the rate of reaction. They remain unchanged and can be used again. They lower the activation energy of the reaction. |
| 7 | Activation energy | Minimum amount of energy required for the reaction to occur |
| 8 | Lock and key model | A model that proposes that each substrate (key) only fits a specific enzyme (lock) |
| 9 | Induced fit model | A model that proposes when the substrate binds with the enzyme, the enzyme changes shape and molds itself to the substrate |


Enzymes (2)

Enzymes catalyse both intracellular and extracellular reactions that determine structures and functions from cellular to whole organism level. There are several factors that affect enzyme activity;


1 Temperature



2 pH




3 Substrate concentration




Enzymes (3)

Enzyme inhibitor- a molecule that binds to an enzyme to decrease its activity. There are two types; competitive and non-competitive inhibitors (see key vocab)


1 Competitive inhibition



2 Non- competitive inhibition




3



ATP

See key vocab for definitions


1



2

| | | |
|-----------------|-----------------|-----------------|
| | ATP to ADP+Pi | ADP+Pi to ATP |
| Reaction | Hydrolysis | Condensation |
| Enzyme involved | ATP hydrolase | ATP synthase |
| Energy profile | Releases energy | Requires energy |

4



Key Vocabulary



| | | |
|---|---------------------------|--|
| 1 | Competitive inhibitor | An inhibitor that resembles the normal substrate binds to the enzyme, usually at the active site, and prevents the substrate from binding. |
| 2 | Non-competitive inhibitor | An inhibitor that does not compete for the active site with the substrate however binds at another place (allosteric site) and changes the shape of the active site so prevents the original substrate binding |
| 3 | ATP | Adenosine triphosphate- An organic compound that provides energy. |
| 4 | ADP + Pi | Adenosine diphosphate + phosphate |
| 5 | Inorganic ions | Atoms or molecules with an electric charge, containing no carbon. Occur in solution in the cytoplasm and body fluids of organisms in high/low concentrations |
| 6 | Cations | Positively charged ions |
| 7 | Anions | Negatively charged ions |

Keywords in key vocab

| | | |
|---|--------------|---|
| 1 | Hydrogen (H) | Determine the pH of body fluids. Higher concentration = lower pH |
| 2 | Iron (Fe) | Part of the prosthetic group in haemoglobin and binds to oxygen |
| 3 | Sodium | Used in co-transport of glucose and amino acids across cell membranes |
| 4 | Phosphate | Essential component of DNA, RNA, ATP |

DNA and RNA

1 DNA and RNA are both polynucleotides



| | | DNA | RNA |
|---|-------------------|--------------------------------------|--|
| 4 | Number of Strands | Two antiparallel strands | One strand |
| 5 | Length | Very long | Relatively short |
| 6 | Pentose Sugar | Deoxyribose | Ribose |
| 7 | Nitrogenous Bases | Adenine, Cytosine, Guanine & Thymine | Adenine, Cytosine, Guanine & Uracil |
| 8 | Function | Store genetic information | Transfer genetic information & forms ribosomes with proteins |

RNA

1 There are 3 types of RNA;

2 Messenger RNA (mRNA)-a transcript copy of a gene which encodes a specific polypeptide

The diagram shows a single-stranded messenger RNA molecule with various colored blocks representing nucleotides.

3 Transfer RNA (tRNA)-carries the polypeptide subunits (amino acids) to the organelle responsible for synthesis (ribosome)

The diagram shows a transfer RNA molecule with a cloverleaf structure and a pink oval representing the ribosome.

4 Ribosomal RNA (rRNA)-a primary component of the ribosome and is responsible for its catalytic activity

The diagram shows a ribosomal RNA molecule with a complex, folded structure and a pink oval representing the ribosome.

DNA replication

| | | |
|---|------------------|--|
| 1 | Polynucleotides | Polymers made up of many nucleotide monomers joined by phosphodiester bonds in a series of condensation reactions |
| 2 | DNA double helix | Held together by hydrogen bonds between complementary base pairs. Adenine + thymine- 2 bonds Cysteine + guanine- 3 H bonds |

3 Semi conservative replication

1- DNA helicase breaks H bonds between the two strands
2-Free nucleotides complementary base pair to the exposed strands
3-DNA polymerase catalyses condensation reactions to join adjacent nucleotides, forming phosphodiester bonds

The diagram illustrates the process of DNA semi-conservative replication. It shows a replication fork where the original (template) DNA strands are being unwound. New DNA strands are synthesized using the template strands. The leading strand is synthesized continuously, while the lagging strand is synthesized discontinuously as Okazaki fragments. DNA polymerase is shown adding free nucleotides to the growing strands. A legend identifies the bases: Adenine (red), Thymine (orange), Cytosine (green), and Guanine (blue).

Key Vocabulary

| | | |
|---|-------------------------------|---|
| 1 | DNA (Deoxyribonucleic acid) | A nucleic acid, composed of nucleotides, that carries genetic instructions. It is double stranded and forms a double helix structure, composed of two polynucleotide chains that interact to form a coil. |
| 2 | RNA (Ribonucleic acid) | A nucleic acid molecule essential in various biological roles in coding, decoding, regulation and expression of genes. |
| 3 | Nucleotide | A structural component of DNA and RNA. Consists of a phosphate group, pentose sugar and organic base |
| 4 | Semi conservative replication | Method by which DNA replicates to form two identical molecules of DNA (consist of one original DNA strand and one newly synthesised DNA strand) |
| 5 | Polarity (water) | A molecule containing polar bonds due to the difference in electronegativity. Consists of one negatively charged and one positively charged end., Eg: Water |
| 6 | Water | Consist of two hydrogen (positively charged) covalently bonded to one oxygen (negatively charged). This causes attraction (hydrogen bonds) between one water molecule and another. |

Water

| | Property | Why is it useful? |
|---|---------------------------------|---|
| 1 | Liquid medium | Provides aquatic habitats, medium for chemical reactions & used for transport |
| 2 | Metabolite | Use in hydrolysis & condensation reactions |
| 3 | High specific heat capacity | Keeps aquatic & cellular environments stable |
| 4 | High latent heat of evaporation | Evaporation has a cooling effect on organisms |
| 5 | Cohesion | Water is 'sticky'- helps to move/pull up the xylem |
| 6 | Surface tension | Allows pond skaters to move on surface |
| 7 | Solvent + transport medium | Dissolves ionic & polar molecules so they can be transported |
| 8 | Reaction medium | Cytoplasm in cells is aqueous solution |
| 9 | Incompressible | Prevents plants from wilting & acts as a hydrostatic skeleton for invertebrates |