
















<div>  <div> <div>Subject: Biology</div> <div>Topic: Organisms response to changes in their internal and external environment</div> <div>Year Group: 13</div> </div> <div>  </div> </div>		
Simple responses – aid survival of a species		
1	Taxis	A simple response whose direction is determined by the direction of the stimulus
2	Kinesis	A non-directional response to a stimulus which brings about increased random movements
3	Tropism	A growth movement of a part of a plant in response to a directional stimulus - Indoleacetic acid (IAA) causes elongation of shoot cells, while it also inhibits root cell elongation in order to cause positive geotropism (to gravity) & phototropism (to light)
4	Reflexes	Rapid responses that don't require conscious thought. Reflexes can quickly protect the body from harm, as it does not involve many synapses, they use simple mechanisms and are localized to the part of the body where they occur.
Receptors		
1	Pacianin corpuscle	Detect changes in pressure in the skin. Increases in pressure cause a deformation of the concentric rings of the Pacinian corpuscle, opening stretch-mediated sodium channels in the membrane. Sodium ions enter the sensory neuron, causing a generator potential which can trigger an action potential.
2	The retina	Contain photoreceptors known as rod cells and cone cells.
3	Rod cells	Detect light in the middle of the visible spectrum. High visual sensitivity. Use the pigment rhodopsin. More abundant than cone cells. Located on the edge of the retina (not in fovea). Show retinal convergence. Provide poor visual acuity.
4	Cone cells	Three different types (detect red, blue and green light) to give colour vision. Use the pigment iodopsin. Found at the fovea. Only one cone to each bipolar cell. Have good visual acuity.
Control of heart rate		
1	Heart is myogenic	Contraction is initiated from within the muscle itself (not from nervous system).
2	SAN → AVN → Bundle of His	The sinoatrial node sends out waves of electrical activity to the left & right atrial wall causing contraction. waves are then passed onto the atrioventricular node, then to the bundle of His, with a slight delay. The bundle of His splits into the Purkynge tissue, causing contraction of the left & right ventricles from the bottom up.
3	Medulla oblongata	Found in the autonomic nervous system – controls impulses to SAN.
4	Receptors – in aorta or carotid arteries	Baroreceptors detect changes in blood pressure, chemoreceptors detect pH changes (e.g due to high CO <sub>2</sub> )
Key Vocabulary		
1	Stimulus	A detectable change in the internal or external environment of an organism
2	Response	The result of a stimulus on an organism
3	Receptor	A cell or organ which detects a stimulus
4	Effector	A cell, tissue, organ or system which responds to a stimulus
5	Coordinator	The link between a sensory neurone and motor neurone in the spinal cord
6	Autonomic	A division of the motor nervous system which carries nerve impulses to glands, smooth and cardiac muscle and is not under voluntary control (subconscious)
7	Sympathetic	An autonomic pathway which stimulates effectors and so speeds up an activity (e.g. heart rate) using noradrenaline
8	Parasympathetic	An autonomic pathway which inhibits effectors and so slows down an activity using acetylcholine
9	Retinal convergence	Many rods connected to one bipolar cell
10	Visual acuity	Ability to resolve fine detail
11	Visual sensitivity	Ability to detect low light intensity
12	Fovea	The point receives the highest intensity of light because light focuses on it by the lens.
13	Blind spot	The point of entry of the optic nerve on the retina which is insensitive to light.
Required Practical 10		
1	Investigation into the effect of an environmental variable e.g. light on the movement of an animal using either a choice chamber or a maze.	

		Subject: Biology	Topic: Organisms response to changes in their internal and external environment		Year Group: 13				
Synapses – gap between 2 neurones			Motor neurones and myelination			Key Vocabulary			
1	Cholinergic synapse	Uses acetylcholine as its neurotransmitter.		1	Structure	Cell body – contains nucleus, cytoplasm and the cell membrane forms dendrites which connect to other neurones Axon – long extension of neurone Schwann cells – surround the axon to form the myelin sheath. Nodes of ranvier – gaps in myelin sheath Axon terminals – connect neurones	1	Resting potential	A potential difference of -65mV found inside a resting neurone relative to its outside, which results in the axon being polarised.
2	Transmission across a synapse	An action potential arrives at the pre-synaptic knob, depolarising the membrane and causes voltage-gated calcium ion channels to open. The influx of Ca <sup>2+</sup> ions causes the synaptic vesicles to fuse with the membrane, releasing the neurotransmitter into the synaptic cleft. The neurotransmitter diffuses and binds receptors on the post synaptic membrane, causing an action potential.					2	Generator potential	A nervous impulse produced by a sensory receptor following transduction (or conversion) of one form of energy into electrical energy
3	Acetylcholinesterase	Breaks down acetylcholine in the cleft. This allows it to be recycled and reused.					3	Threshold value	The minimum level of stimulus needed to trigger an action potential
4	Excitatory or inhibitory synapses	If the neurotransmitter opens Na <sup>+</sup> channels (excitatory) or if it opens chloride or potassium channels causing hyperpolarization (inhibitory)					4	Polarised	Condition used to describe the axon when the inside of an axon is negatively charged relative to the outside (at the resting potential usually around -65mV).
Action potentials			2	Importance of myelination	Increases conduction speed, as well as wider axons or warmer temperatures.		5	Voltage gated channels	Channels in the axon membrane which change shape, and hence open or close, depending on the voltage across the membrane.
3			3	Saltatory conduction	Action potentials only occur at Nodes so faster transmission.		6	Depolarised	Condition used to describe when the inside of the membrane has a positive charge of around +40mV (when an action potential is happening).
1	Definition	A temporary reversal of the charges across the axon membrane which increase from -65mV to +40mV, depolarising the membrane		7	Hyperpolarisation	When the inside of the axon is more negative (relative to the outside) than the usual.			
2	Process	When the neurone receives an impulse from receptors, sodium channels on the dendrites open, leading to the movement of Na <sup>+</sup> ions into the cell causing depolarisation. If this depolarisation reaches the threshold potential it activates voltage-gated sodium channels causing an action potential. After voltage-gated sodium ion channels close, and voltage-gated potassium channels open, repolarization occurs as K <sup>+</sup> ions leave the cell. Outward diffusion of K <sup>+</sup> ions causes hyperpolarisation and the voltage-gated potassium channels close. Finally, the Sodium-potassium pump returns the cell to the resting membrane potential.		8	Repolarisation	When the resting potential of -65mV is re-established the axon is described as this.			
3	Refractory period	Time period after an action potential when it is impossible for a further action potential to be generated because inward movement of sodium ions is prevented because the sodium voltage-gated channels are closed.							
4	All-or-nothing principle	An action potential is exactly the same size, regardless of the size of the stimulus, providing it reaches the threshold value.							

 Beckfoot		Subject: Biology	Topic: Organisms response to changes in their internal and external environment		Year Group: 13					
Neuromuscular junction (NJ) and the Sarcomere				Types of skeletal muscle		Key Vocabulary				
1	Synapse vs neuromuscular junction		<ul style="list-style-type: none"><li>- NJ can only be excitatory</li><li>- NJ link a neurone to a muscle rather than to another neurone</li><li>- Action potential ends at NJ</li><li>- Only motor neurones used</li><li>- NJ always use acetylcholine</li></ul>	1	Slow twitch	<ul style="list-style-type: none"><li>• Used for endurance</li><li>• Contract slowly and for longer</li><li>• Fatigue slowly</li><li>• Carry out aerobic respiration</li><li>• Have lots of mitochondria</li><li>• Lots of myoglobin</li><li>• Small glycogen/phosphocreatine stores</li><li>• Dark colour</li></ul>	1	Myofibrils (muscle fibres)	A microscopic muscle fibre containing sarcoplasm and showing striped isotropic and anisotropic bands of actin and myosin.	
2	Sarcolemma	This is the name of the membrane on the neuromuscular junction with the receptors for the acetylcholine.				2	A Band	Anisotropic bands (dark) where thick and thin filaments overlap.		
3	Sarcomeres			2	Fast twitch	<ul style="list-style-type: none"><li>• Used for short bursts of activity.</li><li>• Light colour</li><li>• In all other aspects they are exactly the <u>opposite</u> to slow twitch</li></ul>	3	I Band	Isotropic bands (light) where thick and thin filaments do not overlap.	
	What are they?	Muscles contain fibres called myofibrils. Each myofibril is made of lots of smaller contracting units called sarcomeres.				4	H zone	The centre of each A-band where there is a lighter-coloured region		
4	The protein filaments in the sarcomere	<p>Actin – thin filaments which “pull” along myosin filaments in contraction</p> <p>Myosin – has “heads” which attach to actin during contraction.</p> <p>Troponin/tropomyosin – wrapped around actin and move when bound Ca<sup>2+</sup> ions to allow myosin to bind.</p>		Contraction of skeletal muscle – sliding filament theory				5	Z line	The centre of each I-band.
				1		How is it initiated?	An action potential travels into the muscle fibre via T tubules, causing release of calcium ions from the sarcoplasmic reticulum. The calcium ions bind to troponin on tropomyosin molecules and cause them to move, exposing the myosin binding site on the actin filament. Myosin attaches to actin forming a actin-myosin cross-bridge.			
				2		Why do muscles shorten during contraction?	ATP hydrolyses to detach the myosin head, allowing reattachment at a further site. This cycle continues, causing sarcomeres to shorten.			
						What happens when nervous stimulation stops?	Ca <sup>2+</sup> ions are actively transported back into the sarcoplasmic reticulum. This allows tropomyosin to block the myosin binding site and muscle contraction stops. ATP can be generated via aerobic or anaerobic respiration. Phosphocreatine generates ATP quickly by adding phosphate to ADP released by the contracting muscle			

		Subject: Biology	Topic: Organisms response to changes in their internal and external environment		Year Group: 13			
Homeostasis and feedback			Control of blood glucose			Key Vocabulary		
1	Definition	Maintenance of a constant internal environment despite internal/external changes.	1	Important to maintain because...	It allows cells to have access to the substrate for respiration, whilst preventing cell damage by dehydration caused by high concentrations.	1	Gluconeogenesis	Happens in the liver - conversion of non-carbohydrates (e.g. glycerol and amino acids) into glucose; literally 'glucose-new-manufacture'. Happens when glycogen supply is exhausted.
2	pH and temperature	Important to regulate to allow optimum enzyme activity and rate of metabolic reactions.	2	High blood glucose	Insulin is released from $\beta$ -cells in the pancreas in order to lower the concentration back to its optimum via negative feedback.	2	Glycogenolysis	Happens in the liver - breakdown of glycogen to glucose – 'breaking down glycogen'. Happens when blood glucose level is lower than normal.
3	Water potential	Important to regulate to prevent cells bursting or shrinking by osmosis.	3	What is the process it uses?	Insulin binds to an insulin receptor, vesicles of glucose transporters fuse with the plasma membrane to allow more glucose to enter the cell. The cell also uses more glucose in respiration and activated enzymes carry out glycogenesis.		Glycogenesis	Happens in the liver - conversion of glucose into glycogen – 'making glycogen'. Happens when blood glucose level is higher than normal.
4	Two types of feedback from change	Positive and negative (see key vocabulary). Predominantly homeostasis is maintained by negative feedback.	4	Low blood glucose	Glucagon is released from $\alpha$ -cells in the pancreas in order to increase the concentration back to its optimum.		4	Negative feedback
Control of blood water potential and osmoregulation			5	What is the process it uses?	Activating enzymes which carry out (glycogenesis) as well as producing glucose from other molecules. Also by Activating enzymes that carry out gluconeogenesis (see key vocabulary).	5	Positive feedback	When the feedback causes the corrective measures to be turned 'on', so makes the system deviate even further from its original or (normal) level
1	Kidney structure – gross (overall) and fine (functional sections called nephrons)		6	Diabetes	Type 1 (insulin not released) and treatment is insulin injection. Type 2 (receptors unresponsive to insulin) and treatment is specialised diet and lifestyle.	6	Osmoregulation	The homeostatic control of the water potential of the blood.
2	Nephrons and the formation of glomerular filtrate	Filtering out various substances in glomerulus due to the smaller diameter of afferent than efferent arteriole. Creates hydrostatic pressure.	7	Adrenaline	Released by the adrenal glands in times of stress and increases blood glucose in anticipation of increased activity. Adrenaline binds to receptors which activates adenylyl cyclase. This converts ATP into cAMP, which acts as a second messenger to activate protein kinase for glycogenolysis.			
3	Reabsorption	The filtrate passes into the Bowman's capsule and where certain ions (in PCT) and water (in DCT) are reabsorbed into the blood whilst the remaining filtrate is excreted as urine.						
4	Maintaining conc. gradient	$\text{Na}^{2+}$ is actively transported out of the PCT into blood to allow glucose & amino acids to co-transport out as $\text{Na}^{2+}$ diffuses into the epithelial cells.						